

MATH 261 EXAM 1

- 1) Find the velocity and acceleration of a particle whose position at time t is given by $\mathbf{r}(t) = (\sin t, \cos t)$. Draw the curve and the vectors.
 - 2) Find $\lim_{(x,y) \rightarrow (0,0)} \frac{3x^2y}{x^2+y^2}$ if it exists. If I define $f(x, y) = \frac{3x^2y}{x^2+y^2}$ for $(x, y) \neq (0, 0)$, can f be extended to a continuous function defined on all of \mathbb{R}^2 ? How?
 - 3) For $f(x, y) = \frac{3x^2y}{x^2+y^2}$, compute the first partial derivatives away from zero.
 - 4) For $z = f(x, y) = \frac{3x^2y}{x^2+y^2}$, find the equation of the tangent plane over the point $(1, 1)$.
 - 5) At $(1, 1)$, in what direction is the function $f(x, y) = \frac{3x^2y}{x^2+y^2}$ increasing fastest?
 - 6) Using the chain rule compute $\frac{d}{dt}(f \circ \mathbf{r})(t)$.
 - 7) Set up the integral giving the volume of the region bounded by the planes $x = 0$, $y = 0$, $z = 0$, $x + y + 2z = 1$.
 - 8) Compute the volume under the surface $z = 1 - x^2 - y^2$ and over the xy -plane.
- Bonus:* Set up any integral giving the volume of the solid above the cone $z = \sqrt{x^2 + y^2}$ and under the sphere $x^2 + y^2 + z^2 = 1$.