

## MATH 396 FINAL EXAM

Notes:

- Please only set up the solutions to the problems. No decimal approximations.
- I have not included any tables—use the usual notation to describe percentiles. E.g.  $F_{p,m,n}$  denotes the 100pth percentile of the  $F$  distribution with  $m$  numerator degrees of freedom and  $n$  denominator degrees of freedom. For normal distribution  $z_p$ , for  $T$  use  $t_{p,n}$ ,  $\chi_{p,n}^2$  for chi-squared, etc.

1) Suppose 100 fair dice are rolled. What is the probability that the sum of the faces showing,  $S$ , equals 100? How about  $200 \leq S \leq 300$ ?

2) Recall that in the derivation of the  $F$ -distribution we computed the *cdf*  $F_{U/V}(t) = P(U \leq tV) = \iint_R f_U f_V$  where  $R = \{(u, v) | u \leq tv\} \subset \mathbb{R}_+^2$ . Find the corresponding region  $R$  for  $F_{UV}$ . Assuming  $U$  and  $V$  are independent random variables from a uniform distribution  $f_U(t) = f_V(t) = 1/\theta$ ,  $t \in [0, \theta]$ , compute  $f_{UV}$  (set up the integrals—*Ans*:  $f_{UV}(t) = \frac{\ln(\theta^2/t)}{\theta^2}$ ,  $t \in [0, \theta^2]$ ).

3) Use the method of moments to estimate the parameter  $\alpha$  in  $f_{UV}(t) = \frac{\ln(\alpha/t)}{\alpha}$ ,  $t \in [0, \alpha]$ . Is the estimator biased? Compute  $\text{Var}(UV)$  and the efficiency of the estimator above. You may find the following integrals useful:  $\int_0^\alpha t^n \ln(\alpha/t) dt = \frac{\alpha^{n+1}}{(n+1)^2}$ .

4) Find the maximum likelihood estimate for the parameter  $p$  in the *pdf*

$$p_X(k) = (1-p)^{k-1}p \quad k = 1, 2, \dots$$

Is this estimator biased?

5) For data  $(y_k)_{k=1}^{10}$  satisfying  $\sum y_k = 1100$  and  $\sum y_k^2 = 130000$  construct a 90% confidence interval for the mean.

6) Let  $U$  and  $V$  be independent  $\chi^2$  random variables with 7 and 9 degrees of freedom, respectively. Find the probability that  $\frac{U/7}{V/9} \in (2.5, 3.3)$ .